

# HEARTBREAKERS™ VITA RED F1

pot tomato



Heartbreakers™ Vita Red F1 is best suited for a 6"/15 cm pot size for retail. The consumers buying such usually re-plant them on their balcony and terrace, and also in their garden. Vita Red was bred for regrowth after harvest of the decorative fruits. Disease tolerance and sturdy leaves makes them attractive. The plant structure is upright with long bunches and bright green fruits for high contrast of the fruit and leaves. Fruit will turn into a glossy sweet red.



## FEATURES

- 6-7" pot size
- Regrowth after first harvest
- Light green fruit for contrast between leaves and fruit

## CULTURAL SHEET

### Properties

Variety number/name	410-080 Vita Red F1
Series name	Heartbreakers™
Species	Lycopersicon esculentum
Common name	(Pot/Determinate) Tomato
Family	Solanum
Type	Annual
Seed weight	1.8-3.0 gram / 1000 sds depending on seed lot and variety
Average germination	85-95%
Plant Height	13.5 - 17.5 inches
Fruit Weight	8-15 gr/fruit depending on culture
Use	<ul style="list-style-type: none"> <li>• Compact snack tomato for indoor use</li> <li>• Compact snack tomato for outdoor use in patio and balcony pots/baskets</li> <li>• Compact snack tomato for kids garden</li> </ul>



1" plug

# HEARTBREAKERS™ VITA RED F1

pot tomato

www.purelineseed.com

222 Wisconsin Dr.  
Jefferson, WI 53549

## YOUNG PLANT CULTURE

### number of seeds/plug

1 for plug size 0.5-1.2 inch

### germination days

1-2 days\* (biological disinfected seeds can take 2 days more)

### germination temp.

64°-70°F (18°-21°C) Covered and high humidity, no light needed

### growing days

14-21 days

### growing temp.

61°-70°F (16°-21°C)

### minimum growing temp.

61°F (16°C) This lengthens the growing days period

### maximum growing temp.

95°F (35°C) This shortens the growing days period, encourage stretching internodes

### optimal day/night temp.

day: 70°F (21°C)

night: 64°F (18°C)

### soil for sowing

Sowing soil with good drainage, EC 1.5. PH 5.8-6.5

### sowing covering

Vermiculite / soil with open structure /app. 2-3 mm thick

### fertilization (f) in the plug

2.5 EC with each watering, NPK 15-10-15 and micro elements

### ready to transplant

Full rooted plug with short internodes. Small young flower could be visible.

## CULTIVATION TIPS DURING YOUNG PLANT GROWING

- Reduce the humidity soon after 70% germination. This prevents stretching of the hypocotyl. For pot tomatoes stretching is not desired.
- The 1-2 week period after cotyledon expansion is the temperature sensitive period, which defines when the first bunch shows. During this period low night temperature exposure 50-60°F of seedlings, in contrast to day temperatures at 64°-70°F promotes the initiation and number of flowers (bunches) on the plant, while also reducing the internode length and the number of leaves preceding the first flower bunch.

## FINISH PLANT CULTURE

### potting soil

Standard soil with good drainage and water storage capabilities EC 2.5 PH 5.8-6.5

### pot size

6-7" optimal 6"

### plugs per pot

1 plug for pots.

### indoor

- Final distance indoor 10" x 10" (16 plts/ m2) for pots

### spacing indoor

Space the plants when the leaves are reaching each other

### outdoor in containers

Planting distance 12" x 12"

### planting soil outdoor

Standard soil with good drainage and water storage capabilities EC 1.5. PH 5.8-6.5

### minimum growing temp.

61°F (18°C) This lengthens the growing days period

### ideal growing temp.

70°-77°F (21°C-25°C)

### optimal day/night temp.

day: 77°F (23°C)

night: 64°F (18°C)

### frost

Plants cannot stand frost

### watering

- Regularly for continues growth, keep the soil moist
- Watering with minimal. 2.5 EC keeps tomatoes healthy

### crop time to saleable product

10-12 weeks after planting, the first bunch will start to show color



- Long days (16 hrs) under relative low light densities increases the dry weight production to 100%. Compared to short day (8 hrs) after 6 weeks from sowing

## CULTIVATION TIPS DURING FINISH PLANT GROWING

- Plants are bred for showing off fruits with low maintenance. They produce long bunches with 14-183 fruits per bunch.
- Put 2-3 sticks around the central stem to keep the plant in balance when fruits are growing.
- High temperatures (80.5°F average D/N) induce pollen infertility/no fruit set. Within the 70-80.5°F range a diurnal fluctuation of at least 5-6°F promotes good pollen fertility.
- Night temperatures of between 60-68°C are ideal. Temperatures below 55°C severely affect the pollination of most cultivars.
- Insects, especially bees and bumble bees support fruit set on tomatoes.
- Tomato plants have a high fertilization need. When the EC is too low, the leaves can turn yellow when the fruits are coloring. This also reduces the taste of the fruit.
- When the flowers show, increase the EC to 3-4, to keep the fertilization in the pot high enough. (EC in the pot can go up to 7-9).
- Tomato plants/leaves can be made sturdier by spraying (MgSO4 -bitter salt and Dipotassium-sulphite (K2SO3)) solutions on the plants (possible combined with other chemicals which need to be used). This has a positive effect on the leaf size and color
- Clay in the soil will stabilize fertilization variation and reduce stretching. 2-5% is advisable, can be increased to 10%.
- Potassium-phosphate (MKPO3) in a concentration of 0.1% improves the quality of tomato plants. This is a good method against phytophthora. For all tomatoes this can lead to a significant reduction of failures.

PureLine Seeds, Inc.